Synergistic antimicrobial activity of apigenin against oral **Pathogens** Su-Mi Cha¹, Gi-Ug Kim², Jeong-Dan Cha^{3*}

¹Department of Oral Microbiology and Institute of Oral Bioscience, Chonbuk National University, Jeonju, South Korea ²Department of Dental Hygiene, Pohang College, Pohang, South Korea ³Institute of Jinan Red Ginseng, Jinan-gun, South Korea

Abstract— Although a broad range of biological and pharmacological activities of apigenin have been reported, the mechanism(s) behind its antibacterial effects are not fully understood. In this study, we investigated the synergistic antibacterial activity of apigenin in combination with existing antimicrobial agents against oral bacteria. The combination effect of apigenin was evaluated against oral bacteria, either alone or with antibiotics, via broth dilution method and checkerboard and time kill assay. MIC/MBC values for apigenin, ampicillin, gentamicin, erythromycin, and vancomycin against all the tested bacteria ranged between 50-200/100-800 microg/mL, 0.0313-16/0.125-32 microg/mL, 2-256/4-512 microg/mL, 0.008-32/0.016-64 microg/mL, and 0.25-64/1-128 microg/mL, respectively. Checkerboard assay revealed synergistic activity in the combination of apigenin with antibiotics at fractional inhibitory concentration index (FICI) <0.5. 1-6 hours of treatment with 1/2 MIC of apigenin with 1/2 MIC of antibiotics resulted from an increase of the rate of killing in units of CFU/mL to a greater degree than was observed with alone. These results suggest that the apigenin is important in the antibacterial actions of oral pathogen agents.

Keywords— Apigenin, antibacterial activity, oral pathogen bacteria, Synergistic effect, Minimum inhibitory concentrations (MICs), Minimum bactericidal concentrations (MBCs)

I. **INTRODUCTION**

Oral diseases are major health problems with dental caries and periodontal diseases among the most important preventable global infectious diseases [1, 2]. Oral health influences the general quality of life and poor oral health is linked to chronic conditions and systemic diseases [3, 4]. The association between oral diseases and the oral microbiota is well established [5]. Of the more than 750 species of bacteria that inhabit the oral cavity, a number are implicated in oral diseases [6-8]. The development of dental caries involves acidogenic and aciduric gram-positive bacteria, primarily the mutans streptococci (Streptococcus mutans and S. sobrinus), lactobacilli, and actinomycetes, which metabolize sucrose to organic acids (mainly lactic acid) that dissolve the calcium phosphate in teeth, causing decalcification and eventual decay [7, 9]. In contrast, periodontal diseases are subgingival conditions that have been linked to anaerobic gram-negative bacteria such as Porphyromonas gingivalis, Actinobacillus sp., Prevotella sp., and Fusobacterium sp. [10, 11]. In periodontal diseases, the areas at or below the gingival crevice become infected causing a cellular inflammatory response of the gingiva and surrounding connective tissue [12, 13]. These inflammatory responses can manifest as gingivitis (extremely common and seen as bleeding of the gingival or gum tissues) or periodontitis (the inflammatory response results in loss of collagen attachment of the tooth to the bone and in loss of bone) [13, 14].

Many plant-derived medicines used in traditional medicinal systems have been recorded in pharmacopeias as agents used to treat infections and a number of these have been recently investigated for their efficacy against oral microbial pathogens [15-19]. Flavonoids have also been shown to exhibit broader bioactivities such as protection of vascular integrity, antihepatotoxicity, anti-inflammatory activity, antitumor effect, antiallergic properties, and antimicrobial effects [20-22]. Apigenin (4,5,7-trihydroxyflavone), a flavone subclass of flavonoid widely distributed in many herbs, fruits, and vegetables, is a substantial component of the human diet and has been shown to possess a variety of biological characteristics, including anticancer, antibacterial, antioxidant, anti-apoptosis, and anti-inflammatory [23-26]. The low antibacterial activity of apigenin alone in Gram-negative bacteria may be due to the presence of lipopolysaccharide and protein-rich outer membrane which covers and protects the internal peptidoglycan wall [27]. The in vitro anti-inflammatory effect of apigenin was studied in many cases. The apigenin decreased cytokines, TNF- α , IL-1 β , and IL-6 in LPS-stimulated human peripheral blood mononuclear cells [28]. Apigenin has been reported to suppress cell proliferation in various cell types, and such an antiproliferative effect has been shown to be associated with PI3K/Akt pathway [29, 30]. Although a broad range of biological and pharmacological activities of apigenin have been reported, the mechanism(s) behind its antibacterial effects are not fully understood. In this study, we investigated the synergistic antibacterial activity of apigenin in combination with existing antimicrobial agents against oral bacteria.

II. MATERIALS AND METHODS

2.1 Bacterial strains

The oral bacterial strains used in this study were: *Streptococcus mutans* ATCC 25175, *Streptococcus sanguinis* ATCC 10556, *Streptococcus sobrinus* ATCC 27607, *Streptococcus ratti* KCTC (Korean collection for type cultures) 3294, *Streptococcus criceti* KCTC 3292, *Streptococcus anginosus* ATCC 31412, *Streptococcus gordonii* ATCC 10558, *Actinobacillus actinomycetemcomitans* ATCC 43717, *Fusobacterium nucleatum* ATCC 10953, *Prevotella intermedia* ATCC 25611, and *Porphylomonas gingivalis* ATCC 33277. Brain-Heart Infusion (Difco Laboratories, Detroit, MI) broth supplemented with 1% yeast extract (Difco) was used for all bacterial strains except *P. intermedia* and *P. gingivalis*. For *P. intermedia* and *P. gingivalis*, BHI broth containing hemin 1 µg/mL (Sigma, St. Louis, MO, USA) and menadione 1 µg/mL (Sigma) was used.

2.2 Minimum inhibitory concentrations/minimum bactericidal concentrations assay

The minimum inhibitory concentrations (MICs) were determined for apigenin by the broth dilution method [18], and were carried out in triplicate. The antibacterial activities were examined after incubation at 37°C for 18 h (facultative anaerobic bacteria), for 24 h (microaerophilic bacteria), and for 1-2 days (obligate anaerobic bacteria) under anaerobic conditions. MICs were determined as the lowest concentration of test samples that resulted in a complete inhibition of visible growth in the broth. MIC₅₀s and MIC₉₀s, defined as MICs at which, 50 and 90%, respectively of oral bacteria were inhibited, were determined. Following anaerobic incubation of **MICs** plates, the minimum bactericidal concentrations (MBCs) were determined on the basis of the lowest concentration of apigenin that kills 99.9% of the test bacteria by plating out onto each appropriate agar plate. Ampicillin, gentamicin, erythromycin, and vancomycin (Sigma) were used as standard antibiotics in order to compare the sensitivity of apigenin against oral bacteria.

2.3 Checker-board dilution test

The antibacterial effects of a combination of apigenin, which exhibited the highest antimicrobial activity, and antibiotics were assessed by the checkerboard test as previously described [18]. The antimicrobial combinations assayed included apigenin with antibiotics, ampicillin, gentamicin, erythromycin, and vancomycin. Serial dilutions of two different antimicrobial agents were mixed in cation-supplemented Mueller-Hinton broth. After 24-48 h of incubation at 37°C, the MICs were determined to be the minimal concentration at which there was no visible growth and MBCs were determined on the basis of the lowest concentration of apigenin that kills 99.9% of the test bacteria by plating out onto each appropriate agar plate. The fractional inhibitory concentration (FIC)/ fractional bactericidal concentration (FBC) index was calculated according to the equation: FIC/FBC index=FIC/FBC_A+FIC/FBC_B=(MIC/MBC of drug A in combination/MIC/MBC of drug A alone)+(MIC/MBC of drug B in combination/MIC/MBC of drug B alone). The FIC and FBC index are the sum of the FICs and FBCs of each of the drugs, which in turn is defined as the MIC and MBC of each drug when it is used in combination divided by the MIC and MBC of the drug when it is used alone. The interaction was defined as synergistic if the FIC and FBC index was less than or equal to 0.5, additive if the FIC and FBC index was greater than 0.5 and less than or equal 1.0, indifferent if the FIC and FBC index was greater than 1.0 and less than or equal to 2.0, and antagonistic if the FIC and FBC index was greater than 2.0.

2.4 Time-kill curves

Bactericidal activities of the drugs under study were also evaluated using time-kill curves on oral bacteria. Tubes containing Mueller-Hinton supplemented to which antibiotics had been added at concentrations of the MIC₅₀ were inoculated with a suspension of the test strain, giving a final bacterial count between $5\sim6.6\times10^6$ CFU/ml. The tubes were thereafter incubated at 37° C in an anaerobic chamber and viable counts were performed at 0, 0.5, 1, 2, 3, 4, 5, 6, 12 and 24 h after addition of antimicrobial agents, on agar plates incubated for up to 48 h in anaerobic chamber at 37° C. Antibiotic carryover was minimized by washings by centrifugation and serial 10-fold dilution in sterile phosphate-buffered saline, pH 7.3. Colony counts were performed in duplicate, and means were taken. The solid media used for colony counts were BHI agar for streptococci and BHI agar containing hemin and menadione for *P. intermedia* and *P. gingivalis*.

III. **RESULTS AND DISCUSSION**

The main etiological factor of dental caries and periodontal disease is dental plaque [31-33]. Therefore, it is reasonable to search for natural products that have antiplaque properties and antimicrobial activity against oral pathogens [16, 19, 34-35]. Apigenin was evaluated for their antimicrobial activities against eleven common bacterial species present in the oral cavity. The results of the antimicrobial activity showed that apigenin exhibited antimicrobial activities against cariogenic bacteria (MICs, 25 to 200 µg/mL; MBCs, 100 to 800 µg/mL), against periodontopathogenic bacteria (MICs, 100 to 200 µg/mL; MBCs, 200 to 400 µg/mL) and for ampicillin, either 0.0313/0.125 or 16/32 µg/mL; for gentamicin, either 2/4 or 256/512 µg/mL; for erythromycin, either 0.008/0.016 or 32/64; for vancomycin, either 0.25/1 or 64/128 on tested all bacteria (Table 1). The MIC₅₀ and MIC₉₀ ranges of apigenin were from 6.25 to 12.5 µg/mL and 25 to 200 µg/mL, respectively. The apigenin showed stronger antimicrobial activity against *S. ratti*, *S. gordonii*, and *P. intermedia* than another bacteria (MIC/MBC, 25/50-100 µg/mL) and the range of MIC₅₀ and MIC₉₀ were 6.25 µg/mL and 25 µg/mL.

TABLE I					
ANTIBACTERIAL ACTIVITY OF APIGENIN AND ANTIBIOTICS IN ORAL BACTERIA					

	Apigenin			Ampicillin	Gentamicin	Erythromycin	Vancomycin
Samples	MIC _{50<}	MIC _{90<}	MIC/MBC		MIC	C/MBC	
S. mutans	25	100	100/200	0.125/0.25	8/16	0.063/0.125	1/2
ATCC 25175 ¹							
S. sanguinis	12.5	50	50/200	0.25/0.5	16/32	0.016/0.031	0.25/1
ATCC 10556							
S. sobrinus	25	100	100/200	0.0313/0.125	16/32	0.031/0.063	1/2
ATCC 27607							
S. ratti	12.5	50	50/100	0.125/0.5	8/16	0.008/0.016	0.5/1
KCTC 3294 ²							
S. criceti	6.25	25	25/100	0.0313/0.125	8/16	0.125/0.25	1/4
КСТС 3292							
S. anginosus	50	200	200/800	0.0625/0.25	8/16	0.125/0.5	1/4
ATCC 31412							
S. gordonii	12.5	50	50/100	0.0625/0.25	16/32	0.031/0.063	0.5/1
ATCC 10558							
A. actinomycetemcomitans	50	200	200/400	16/32	8/16	0.125/0.25	2/4
ATCC 43717							
F. nucleatum	25	100	100/200	8/16	2/4	32/64	64/128
ATCC 51190							
P. intermedia	50	200	200/400	1/2	32/32	16/32	16/36
ATCC 49049							
P. gingivalis	25	200	200/400	0.5/0.5	256/512	2/8	8/16
ATCC 33277							

¹American Type Culture Collection (ATCC)

Natural products, polyphenols are a major source of chemical diversity and have provided important therapeutic agents for many oral bacterial diseases [36-38]. Combinations of some herbal materials and different antibiotics might affect the inhibitory effect of these antibiotics [18, 39, 40]. The synergistic effects of apigenin alone or with antibiotics were evaluted in oral bacteria (Table 2 and 3). In combination with apigenin, the MIC for ampicillin was reduced \geq 4-fold in tested bacteria, except *S. anginosus*, producing a synergistic effect as defined by FICI \leq 0.5. The MBC for ampicillin was shown synergistic effects in *S. mutans*, *S. sanguinis*, *S. ratti*, *S. gordonii*, *F. nucleatum*, *P. intermedia*, and *P. gingivalis* by FBCI \leq 0.5 (Table 2). In combination with apigenin, the MIC for gentamicin was reduced \geq 4-fold in tested bacteria expect *S. ratti* by FICI \leq 0.5 and MBC in *S. mutans*, *S. sobrinus*, *S. criceti*, *S. anginosus*, *S. gordonii*, *F. nucleatum*, *P. intermedia*, and *P. gingivalis* by FBCI \leq 0.5 (Table 3). Moreover, the MIC for erythromycin with apigenin was reduced \geq 4-fold in tested bacteria, except *S. sanguinis*, *S. rattii*, and *S. anginosus*, producing a synergistic effect as defined by FICI \leq 0.5 (Table 3). Moreover, the MIC for erythromycin with apigenin was reduced \geq 4-fold in tested bacteria, except *S. sanguinis*, *S. rattii*, and *S. anginosus*, producing a synergistic effect as defined by FICI \leq 0.5 and the MBC for erythromycin was shown synergistic effects in *S. mutans*, *S. gordonii*, *A. actinomycetemcomitans*, and *P. gingivalis* by FBCI \leq 0.5 (Table 2).

		MIC/MBC (µg/ml)				
Strains	Agent	Alone	Combination ¹	FIC/FBC	FICI/FBCI ²	Outcome
S. mutans	Apigenin	100/200	12.5/50	0.125/0.125	0.375/0.375	Synergistic/
ATCC 25175 ³	Ampicillin	0.125/0.25	0.0313/0.0625	0.25/0.25		Synergistic
S. sanguinis	Apigenin	50/200	12.5/50	0.25/0.25	0.5/0.5	Synergistic/ Synergistic
ATCC 10556	Ampicillin	0.25/0.5	0.0625/0.125	0.25/0.25		
S. sobrinus	Apigenin	100/200	25/50	0.25/0.25	0.5/0.75	Synergistic/ Additive
ATCC 27607	Ampicillin	0.0625/0.125	0.0156/0.0625	0.25/0.5		
S. ratti	Apigenin	50/100	12.5/25	0.25/0.25	0.5/0.5	Synergistic/ Synergistic
KCTC 3294 ⁴	Ampicillin	0.25/0.5	0.0625/0.125	0.25/0.25		
S. criceti	Apigenin	25/100	6.25/12.5	0.25/0.125	0.5/0.625	Synergistic/ Additive
КСТС 3292	Ampicillin	0.0625/0.125	0.0156/0.0625	0.25/0.5		
S. anginosus	Apigenin	200/800	50/200	0.25/0.25	0.75/0.75	Additive/ Additive
ATCC 31412	Ampicillin	0.125/0.25	0.0625/0.125	0.5/0.5		
S. gordonii	Apigenin	50/100	12.5/12.5	0.25/0.125	0.5/0.375	Synergistic/ Synergistic
ATCC 10558	Ampicillin	0.0625/0.25	0.0156/0.0625	0.25/0.25		
A. actinomycetemcomitans	Apigenin	200/400	25/100	0.125/0.25	0.25/0.75	Synergistic/ Additive
ATCC 43717	Ampicillin	16/32	2/16	0.125/0.5		
F. nucleatum	Apigenin	100/200	25/50	0.25/0.25	0.5/0.5	Synergistic/ Synergistic
ATCC 51190	Ampicillin	16/32	4/8	0.25/0.25		
P. intermedia	Apigenin	200/400	50/100	0.25/0.25	0.375/0.5	Synergistic/ Synergistic
ATCC 49049	Ampicillin	2/4	0.25/1	0.125/0.25		
P. gingivalis	Apigenin	200/400	50/100	0.25/0.25	0.5/0.5	Synergistic/ Synergistic
ATCC 33277	Ampicillin	0.5/1	0.125/0.25	0.25/0.25		

 TABLE 2

 Synergistic effects of apigenin with ampicillin against oral bacteria.

¹The MIC and MBC of apigenin with ampicillin

² The fractional inhibitory concentration index (FIC index)

³American Type Culture Collection (ATCC)

		MIC/MBC (µg/ml)		FIC	FICI ²	Outcome
Strains	Agent	Alone	Combination ¹			
S. mutans	Apigenin	100/200	12.5/50	0.125/0.25	0 375/0 5	Synergistic/ Synergistic
ATCC 25175 ³	Gentamicin	8/16	2/4	0.25/0.25	0.373/0.3	
S. sanguinis	Apigenin	50/200	12.5/50	0.25/0.25	0.5/0.75	Synergistic/
ATCC 10556	Gentamicin	16/32	4/16	0.25/0.5		Additive
S. sobrinus	Apigenin	100/200	12.5/25	0.125/0.125	0 375/0 375	Synergistic/ Synergistic
ATCC 27607	Gentamicin	16/32	4/16	0.25/0.5	-	
S. ratti	Apigenin	50/100	12.5/25	0.5/0.25	0.75/0.75	Additive/ Additive
KCTC 3294 ⁴	Gentamicin	16/16	4/8	0.25/0.5		
S. criceti	Apigenin	25/100	6.25/12.5	0.25/0.125	0.5/0.375	Synergistic/ Synergistic
КСТС 3292	Gentamicin	8/16	2/4	0.25/0.25		
S. anginosus	Apigenin	200/800	50/200	0.25/0.25	0.375/0.5	Synergistic/ Synergistic
ATCC 31412	Gentamicin	16/32	2/8	0.125/0.25		
S. gordonii	Apigenin	50/100	6.25/25	0.125/0.25	0.375/0.5	Synergistic/
ATCC 10558	Gentamicin	16/32	4/8	0.25/0.25	-	Synergistic
A. actinomycetemcomitans	Apigenin	200/400	50/100	0.25/0.25	0.5/0.75	Synergistic/ Additive
ATCC 43717	Gentamicin	8/16	2/8	0.25/0.5		
F. nucleatum	Apigenin	100/200	12.5/25	0.125/0.125	0.375/0.375	Synergistic/ Synergistic
ATCC 51190	Gentamicin	4/8	1/2	0.25/0.25		
P. intermedia	Apigenin	200/400	50/100	0.25/0.25	0.375/0.75	Synergistic/
ATCC 25611	Gentamicin	32/32	4/16	0.125/0.5		Additive
P. gingivalis	Apigenin	200/400	50/100	0.25/0.25	0.375/0.5	Synergistic/
ATCC 33277	Gentamicin	256/512	32/128	0.125/0.25		Synergistic

 TABLE 3
 Synergistic effects of apigenin with gentamicin against oral bacteria.

¹The MIC and MBC of apigenin with gentamicin

² The fractional inhibitory concentration index (FIC index)

³American Type Culture Collection (ATCC)

 TABLE 4

 Synergistic effects of apigenin with erythromycin against oral bacteria

		MIC/MBC (µg/ml)		FIC	FICI ²	Outcome
Strains	Agent	Alone	Combination ¹			outcome
S. mutans	Apigenin	100/200	25/50	0.25/0.25	0.375/0.5	Synergistic/
ATCC 25175 ³	Erythromycin	0.063/0.125	0.008/0.031	0.125/0.25		Synergistic
S. sanguinis	Apigenin	50/200	12.5/50	0.25/0.25	0.75/0.75	Additive/
ATCC 10556	Erythromycin	0.016/0.031	0.008/0.016	0.5/0.5		Additive
S. sobrinus	Apigenin	100/200	25/100	0.25/0.5	0.5/0.75	Synergistic/
ATCC 27607	Erythromycin	0.031/0.063	0.008/0.016	0.25/0.25		Additive
S. ratti	Apigenin	50/100	12.5/25	0.25/0.25	0.75/0.75	Additive/
KCTC 3294⁴	Erythromycin	0.008/0.016	0.004/0.008	0.5/0.5	0.7570.75	Additive
S. criceti	Apigenin	25/100	6.25/25	0.25/0.25	0.5/0.5	Synergistic/
KCTC 3292	Erythromycin	0.125/0.25	0.031/0.063	0.25/0.25		Synergistic
S. anginosus	Apigenin	200/800	50/100	0.25/0.125	0.75/0.375	Additive/
ATCC 31412	Erythromycin	0.125/0.5	0.063/0.0125	0.5/0.25		Synergistic
S. gordonii	Apigenin	50/100	12.5/25	0.25/0.25	0.5/0.5	Synergistic/
ATCC 10558	Erythromycin	0.031/0.063	0.008/0.016	0.25/0.25		Synergistic
A. actinomycetemcomitans	Apigenin	200/400	50/100	0.25/0.25	0.5/0.5	Synergistic/
ATCC 43717	Erythromycin	0.125/0.25	0.031/0.063	0.25/0.25		Synergistic
F. nucleatum	Apigenin	100/200	25/50	0.25/0.25	0.5/0.75	Synergistic/
ATCC 51190	Erythromycin	32/64	8/32	0.25/0.5		Additive
P. intermedia	Apigenin	200/400	50/200	0.25/0.5	0.5/0.75	Synergistic/
ATCC 25611	Erythromycin	16/32	4/8	0.25/0.25]	Additive
P. gingivalis	Apigenin	200/400	50/100	0.25/0.25	0.375/0.5	Synergistic/
ATCC 33277	Erythromycin	2/8	0.25/2	0.125/0.25]	Synergistic

¹The MIC and MBC of apigenin with erythromycin

² The fractional inhibitory concentration index (FIC index)

³American Type Culture Collection (ATCC)

		MIC/MBC (µg/ml)		FIC	FICI ²	Outcome
Strains	Agent	Alone	Combination ¹			Outcome
S. mutans	Apigenin	100/200	25/100	0.25/0.5	0.5/0.75	Synergistic/
ATCC 25175 ³	Vancomycin	1/2	0.25/0.5	0.25/0.25	0.5/0.75	Additive
S. sanguinis	Apigenin	50/200	12.5/50	0.25/0.25	0.5/0.375	Synergistic/
ATCC 10556	Vancomycin	0.25/1	0.063/0.125	0.25/0.125		Synergistic
S. sobrinus	Apigenin	100/200	25/50	0.25/0.25	0.5/0.5	Synergistic/ Synergistic
ATCC 27607	Vancomycin	1/2	0.25/0.5	0.25/0.25	0.5/0.5	
S. ratti	Apigenin	50/100	12.5/50	0.25/0.5	0.5/0.75	Synergistic/
KCTC 3294 ⁴	Vancomycin	0.5/1	0.125/0.25	0.25/0.25	0.5/0.75	Additive
S. criceti	Apigenin	25/100	6.25/25	0.25/0.25	0.5/0.5	Synergistic/
КСТС 3292	Vancomycin	1/4	0.25/1	0.25/0.25		Synergistic
S. anginosus	Apigenin	200/800	50/100	0.25/0.125	0.5/0.25	Synergistic/
ATCC 31412	Vancomycin	1/4	0.25/0.5	0.25/0.125	0.070.20	Synergistic
S. gordonii	Apigenin	50/100	12.5/25	0.25/0.25	0.5/0.5	Synergistic/
ATCC 10558	Vancomycin	0.5/1	0.125/0.25	0.25/0.25		Synergistic
A. actinomycetemcomitans	Apigenin	200/400	50/100	0.25/0.25	0.5/0.75	Synergistic/
ATCC 43717	Vancomycin	2/4	0.5/2	0.25/0.5		Additive
F. nucleatum	Apigenin	100/200	25/100	0.25/0.5	0.5/0.75	Synergistic/
ATCC 51190	Vancomycin	64/128	16/32	0.25/0.25		Additive
P. intermedia	Apigenin	200/400	50/100	0.25/0.25	0.5/0.75	Synergistic/
ATCC 25611	Vancomycin	16/36	4/16	0.25/0.5		Additive
P. gingivalis	Apigenin	200/400	50/100	0.25/0.25	0.5/0.5	Synergistic/
ATCC 33277	Vancomycin	8/16	2/4	0.25/0.25		Synergistic

 TABLE 5

 Synergistic effects of apigenin with vancomycin against oral bacteria.

¹The MIC and MBC of apigenin with vancomycin

² The fractional inhibitory concentration index (FIC index)

³American Type Culture Collection (ATCC)

Flavonoid complexes attach with extra cellular soluble protein and with bacterial cell wall. Thus they exhibit antibacterial activity [41, 42]. Naturally derived apigenin was reported to have potential antimicrobial activity [26, 27, 43]. Trihydroxyflavone or apigenin is a naturally occurring bioflavonoid abundantly present in fruits and vegetables whose antibacterial activity against certain strains of Gram-negative and Gram-positive bacteria like *Escherichia coli, Staphylococcus aureus, Bacillus cereus, and Pseudomonas aeruginosa* has been reported recently [26,27,43,44].

In this study, apigenin, a flavone compound found in several plants also shows susceptibility on gram-positive bacteria as well as gram-negative bacteria. The bacterial effect of apigenin with ampicillin or gentamicin against oral bacteria was confirmed by time-kill curve experiments. The apigenin (MIC or MIC_{50}) alone resulted rate of killing increasing or not changing in CFU/ml at time dependent manner, with a more rapid rate of killing by apigenin (MIC₅₀) with ampicillin or/and gentamicin (MIC₅₀) (Fig 1-2). A strong bactericidal effect was exerted in drug combinations.



FIG. 1. TIME-KILL CURVES OF MIC OF APIGENIN ALONE AND ITS COMBINATION WITH MIC50 OF AMP OR GEN AGAINST. S. MUTANS, S. SANGUINIS, S. SOBRINUS, S. ANGINOSUS, S. CRICETI, AND S. RATTI. BACTERIA WERE INCUBATED WITH APIGENIN (●), APIGENIN + AMP (○), AND APIGENIN + GEN (▼) OVER TIME. DATA POINTS ARE THE MEAN VALUES±S.E.M. OF SIX EXPERIMENTS. CFU, COLONY-FORMING UNITS.



 FIG. 2. TIME-KILL CURVES OF MIC OF APIGENIN ALONE AND ITS COMBINATION WITH MIC50 OF AMP OR GEN AGAINST S. GORDONII, A. ACTINOMYCETEMCOMITANS, F. NUCLEATUM, P. INTERMEDIA, AND P.
 GINGIVALIS. BACTERIA WERE INCUBATED WITH APIGENIN (●), APIGENIN + AMP (○), AND APIGENIN + GEN (▼) OVER TIME. DATA POINTS ARE THE MEAN VALUES±S.E.M. OF SIX EXPERIMENTS. CFU, COLONY-FORMING UNITS.

IV. CONCLUSION

In conclusion, these findings suggest that apigenin, a flavone compound found in several plants fulfills the conditions required of a novel cariogenic bacteria and periodontal pathogens, particularly bacteroides species drug and may be useful in the future in the treatment of oral bacteria.

ACKNOWLEDGEMENTS

This work was supported by a Korea Research Foundation Grant funded by the Korean Government (KRF-20110023479).

CONFLICT OF INTEREST STATEMENT

The authors have declared no conflict of interest.

AUTHORS' CONTRIBUTION

Jeong-Dan Cha has substantial contributions to conception and design and drafting and revising it. Su-Mi Cha and Gi-Ug Kim have substantial contributions to acquisition and analysis of data.

REFERENCES

- P. D. Marsh "Dental plaque as a biofilm and a microbial community implications for health and disease," BMC Oral Health, vol. 15, no. 6, pp. S14, 2006.
- [2] B. Grossner-Schreiber, T. Fetter, J. Hedderich, T. Kocher, S. Schreiber, and S. Jepsen "Prevalence of dental caries and periodontal disease in patients with inflammatory bowel disease: a case-control study," J Clin Periodontol., vol. 33. No.7, pp. 478-484, 2006.
- [3] A. C. Chi, B. W. Neville, J. W. Krayer, and W. C. Gonsalves "Oral manifestations of systemic disease," Am Fam Physician., vol. 82, no. 11, pp. 1381-1318, 2010.
- [4] A. Sheiham, J. G. Steele, W. Marcenes, G. Taskos, S. Finch, and A.W. G. Walls "Prevalence of impacts of dental and oral disorders and their effects on eating among older people: a national survey in Great Britain," Community Dent Oral Epidemiol., vol. 29, no. 3, pp. 195- 203, 2001.
- [5] J.A. Aas, B.J. Paster, L.N. Stokes, I. Olsen, and F.E. Dewhirst "Defining the normal bacterial flora of the oral cavity," J Clin Microbiol., vol. 43, no. 11, pp. 5721-5732, 2005.
- [6] L.M. Collins, and C. Dawes "The surface area of the adult human mouth and thickness of the salivary film covering the teeth and oral mucosa," J Dent Res., vol. 66, no. 8, pp. 1300-1302, 1987.
- [7] P. D. Marsh "Dental plaque as a microbial biofilm," Caries Res., vol. 38, no. 3, pp. 204-211, 2004.
- [8] L. Sbordone, and C. Bortolaia "Oral microbial biofilms and plaque-related diseases: microbial communities and their role in the shift from oral health to disease," Clin Oral Investig, vol. 7, no. 4, pp. 181-188, 2003.
- [9] I. Kleinberg "A mixed-bacteria ecological approach to understanding the role of the oral bacteria in dental caries causation: an alternative to Streptococcus mutans and the specific-plaque hypothesis," Crit Rev Oral Biol Med., vol. 13, no. 2, pp. 108-125, 2002.
- [10] E.A. Palombo "Traditional medicinal plant extracts and natural products with activity against oral bacteria: potential application in the prevention and treatment of oral diseases," Evid based Complement Alt Med., vol. 2011, pp. 680354, 2011.
- [11] C.M. Ardila, M.A. López, and I.C. Guzmán "Positive correlations between presence of gram negative enteric rods and Porphyromonas gingivalis in subgingival plaque," Acta odontol Latinoam., vol. 24, no. 1, pp. 15-19, 2011.
- [12] S. S. Lagdive, S. B. Lagdive, A. Mani, R. Anarthe, G. Pendyala, B. Pawar, and P. P. Marawar "Correlation of mast cells in periodontal disease," J Indian Soc Periodontol., vol. 17, no. 1, pp. 63-67, 2013.
- [13] D.T. Graves, J. Li, and D. L. Cochran "Inflammation and uncoupling as mechanisms of periodontal bone loss," J Dent Res., vol. 90, no. 2, pp. 143-153, 2011.
- [14] D. A. Van Strydonck, D. E. Slot, U. Van der Velden, F. Van der Weijden "Effect of a chlorhexidine mouthrinse on plaque, gingival inflammation and staining in gingivitis patients: a systematic review," J Clin Periodontol., vol. 39, no. 11, pp. 1042-1055, 2012.
- [15] B.B. Mishra, and V. K. Tiwari "Natural products: an evolving role in future drug discovery," Eur J Med Chem., vol. 46, no. 10, pp. 4769-4807, 2011.
- [16] J. S. Choi, Y. M. Ha, C. U. Joo, K. K. Cho, S. J. Kim, and I. S. Choi "Inhibition of oral pathogens and collagenase activity by seaweed extracts," J Environ Biol., vol. 33, no. 1, pp. 115-121, 2012.
- [17] A. Bag, S.K. Bhattacharyya, N. K. Pal, and R.R. Chattopadhyay, 2012. "In vitro antibacterial potential of Eugenia jambolana seed extracts against multidrug-resistant human bacterial pathogens," Microbiol Res., vol. 167, no. 6, pp. 352-357, 2012.
- [18] J. D. Cha, M.R. Jeong, S. I. Jeong, and K. Y. Lee "Antibacterial activity of sophoraflavanone G isolated from the roots of Sophora flavescens." J Microbiol Biotechnol., vol. 17, no. 5, pp. 858-864, 2007.

- [19] A. Vermani, Navneet, and Prabhat "Screening of Quercus infectoria gall extracts as anti-bacterial agents against dental pathogens," Indian J Dent Res., vol. 20, no. 3, pp. 337-339, 2009.
- [20] N. S. Radulović, M. B. Stojković, S. S. Mitić, P. J. Randjelović, I. R. Ilić, N. M. Stojanović, and Z. Z. Stojanović-Radić "Exploitation of the antioxidant potential of Geranium macrorrhizum (Geraniaceae): hepatoprotective and antimicrobial activities," Nat Prod Commun., vol. 7, no. 12, pp. 1609-1614, 2012.
- [21] S. Fu, C. Sun, X. Tao, and Y. Ren "Anti-inflammatory effects of active constituents extracted from Chinese medicinal herbs against Propionibacterium acnes," Nat Prod Res., vol. 26, no. 18, pp. 1746-1749, 2012.
- [22] R. Prasad, and S. K. Katiyar "Bioactive phytochemical proanthocyanidins inhibit growth of head and neck squamous cell carcinoma cells by targeting multiple signaling molecules," PLoS One, vol. 7, no. 9, pp. e46404, 2012.
- [23] H. Wei, L. Tye, E. Bresnick, and D. F. Birt "Inhibitory effect of apigenin, a plant flavonoid, on epidermal ornithine decarboxylase and skin tumor promotion in mice," Cancer Res., vol. 50, no. 3, pp. 499-502, 1990.
- [24] H. Sharma, R. Kanwal, N. Bhaskaran, and S. Gupta "Plant flavone apigenin binds to nucleic acid bases and reduces oxidative DNA damage in prostate epithelial cells," PLoS One, vol. 9, no. 3, pp. e91588, 2014.
- [25] J. Wang, Y.T. Liu, L. Xiao, L. Zhu, Q. Wang, and T. Yan "Anti-Inflammatory Effects of Apigenin in Lipopolysaccharide-Induced Inflammatory in Acute Lung Injury by Suppressing COX-2 and NF-kB Pathway," Inflammation, vol. 37, no. 6, pp. 2085-2090, 2014.
- [26] K. Banerjee, S. Banerjee, S. Das, and M. Mandal "Probing the potential of apigenin liposomes in enhancing bacterial membrane perturbation and integrity loss," J Colloid Interface Sci., vol. 1, no. 453, pp. 48-59, 2015.
- [27] G. Eumkeb, and S. Chukrathok "Synergistic activity and mechanism of action of ceftazidime and apigenin combination against ceftazidime-resistant Enterobacter cloacae," Phytomedicine, vol. 15, no 20, pp. 262-269, 2013.
- [28] S. Hougee, A. Sanders, J. Faber, Y. M. Graus, W. B. van den Berg, J. Garssen, H. F. Smit, and M. A. Hoijer "Decreased proinflammatory cytokine production by LPS-stimulated PBMC upon in vitro incubation with the flavonoids apigenin, luteolin or chrysin, due to selective elimination of monocytes/macrophages," Biochem Pharmacol., vol. 69, no. 2, pp. 241-248, 2005.
- [29] J. Fang, Y. Y. Bao, S. H. Zhou, and J. Fan "Apigenin inhibits the proliferation of adenoid cystic carcinoma via suppression of glucose transporter-1," Mol Med Rep., vol. 12, no. 5, pp. 6461-6466, 2015.
- [30] W.J. Lee, W.K. Chen, C.J. Wang, W.L. Lin, and T.H. Tseng "Apigenin inhibits HGF-promoted invasive growth and metastasis involving blocking PI3K/Akt pathway and beta 4 integrin function in MDA-MB-231 breast cancer cells," Toxicol Appl Pharmacol., vol. 226, no. 2. pp.178-191, 2008.
- [31] E. Pasich, M. Walczewska, A. Pasich, and J. Marcinkiewicz "Mechanism and risk factors of oral biofilm formation," Postepy Hig Med Dosw., vol. 2, no. 67, pp. 736-741, 2013.
- [32] C. J. Seneviratne, C. F. Zhang, and L. P. Samaranayake "Dental plaque biofilm in oral health and disease," Chin J Dent Res., vol. 14, no. 2, pp.87-94, 2011.
- [33] W. Loesche "Dental caries and periodontitis: contrasting two infections that have medical implications," Infect Dis Clin North Am., vol. 21, no. 2, pp. 471-502, 2007.
- [34] K.A. Hussain, B. Tarakji, B.P. Kandy, J. John, J. Mathews, V. Ramphul, and D.D. Divakar "Antimicrobial effects of citrus sinensis peel extracts against periodontopathic bacteria: an in vitro study," Rocz Panstw Zakl Hig., vol. 66, no. 2, pp.173-178, 2015.
- [35] B. Kouidhi, Y. M. Al Qurashi, and K. Chaieb "Drug resistance of bacterial dental biofilm and the potential use of natural compounds as alternative for prevention and treatment," Microb Pathog., vol. 80, no. 21, pp.39-49, 2015.
- [36] J. K. Patra, E. S. Kim, K. Oh, H. J. Kim, Y. Kim, and K. H. Baek "Antibacterial effect of crude extract and metabolites of Phytolacca americana on pathogens responsible for periodontal inflammatory diseases and dental caries," BMC Complement Altern Med., vol. 20, no. 14, pp.343, 2014.
- [37] L. Bonifait, and D. Grenier "Cranberry polyphenols: potential benefits for dental caries and periodontal disease," J Can Dent Assoc., vol. 76, pp. a130, 2010.
- [38] S. Petti, and C. Scully "Polyphenols, oral health and disease: A review," J Dent., vol. 37, no. 6, pp. 413-423, 2009.
- [39] S. Hemaiswarya, A. K. Kruthiventi, and M. Doble "Synergism between natural products and antibiotics against infectious diseases," Phytomedicine, vol. 15, no. 8, pp. 639-652, 2008.
- [40] X.H. Ma, C. J. Zheng, L. Y. Han, B. Xie, J. Jia, Z. W. Cao, Y. X. Li, and Y. Z. Chen "Synergistic therapeutic actions of herbal ingredients and their mechanisms from molecular interaction and network perspectives," Drug Discov Today., vol. 14, no. 11-12, pp. 579-588, 2009.
- [41] J. B. Harborne, and C. A. Williams "Advances in flavonoid research since 1992," Phytochemistry, vol. 55, pp. 481-504, 2000.
- [42] P. D. Stapleton, S. Shah, K. Ehlert, Y. Hara, and P. W. Taylor "The beta-lactam-resistance modifier (-)-epicatechin gallate alters the architecture of the cell wall of Staphylococcus aureus," Microbiology, vol. 153, no. Pt 7, pp. 2093-2103, 2007.
- [43] H. Koo, B. Schobel, K. Scott-Anne, G. Watson, W. H. Bowen, J. A. Cury, P. L. Rosalen, and Y. K. Park "Apigenin and tt-farnesol with fluoride effects on S. mutans biofilms and dental caries," J Dent Res., vol. 84, no. 11, pp. 1016-1020, 2005.
- [44] L. E. Alcaraz, S. E. Blanco, O. N. Puig, F. Tomas, and F. H. Ferretti "Antibacterial activity of flavonoids against methicillin-resistant Staphylococcus aureus strains," J Theor Biol., vol. 205, no. 2, pp. 231-240, 2000.